arXiv:1207.4060v1 [math.SP] 17 Jul 2012

MULTIPLICITY OF THE LOWEST EIGENVALUE OF NON-COMMUTATIVE HARMONIC OSCILLATORS

FUMIO HIROSHIMA AND ITARU SASAKI

Abstract. The multiplicity of the lowest eigenvalue E of the so-called non-commutative harmonic oscillator $Q(\alpha,\beta)$ is studied. It is shown that E is simple for α and β in some region.

1. Definition and main results

Recently a special attention is payed to studying the spectrum of self-adjoint operators with *non-commutative* coefficients. It is considered not only as mathematics but also physics experiments. A historically important model is the Dirac operator, and the Rabi model and the Jaynes-Cumming model are prevalent in cavity QED. See [HH12] and references therein. The non-commutative harmonic oscillator is a quantum system defined by the Hamiltonian:

$$Q = Q(\alpha,\beta) = A \otimes \left(-\frac{1}{2}\frac{d^2}{dx^2} + \frac{1}{2}x^2\right) + J \otimes \left(x\frac{d}{dx} + \frac{1}{2}\right), \tag{1.1}$$

where $A = \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix}$, $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, and $\alpha, \beta > 0$ parameters with $\alpha\beta > 1$. Operator Q defines a positive self-adjoint operator acting in the Hilbert space $\mathcal{H} = \mathbb{C}^2 \otimes L^2(\mathbb{R})$.

The non-commutative harmonic oscillator Q has been introduced by Parmeggiani and Wakayama [PW01, PW02a, PW02b, PW03], and the spectral property of Q is considered in [Par04, Par06, Par08a] from the pseudo-differential-calculus point of view. It can be seen that Q has purely discrete spectrum $\lambda_1 \le \lambda_2 \le \cdots \le \lambda_n \le \cdots \uparrow \infty$, where the eigenvalues are counted with multiplicity. One can define the so-called spectral zeta function associated with Q as

$$\zeta_Q(s) = \sum_{n=1}^{\infty} \frac{1}{\lambda_n^s}.$$

When $\alpha = \beta$, Q is unitarily equivalent to the direct some of harmonic oscillators, and $\lambda_{2m-1} = \lambda_{2m} = \sqrt{\alpha^2 - 1}(m + \frac{1}{2})$, and thus ζ_Q with $\alpha \ne \beta$ can be regarded as a q-deform of the Riemann zeta-function. Analytic properties of the spectral zeta-function

---

Date: July 18, 2012.

2000 *Mathematics Subject Classification.* 35P05, 35P15.

Key words and phrases. non-commutative harmonic oscillator, multiplicity, lowest eigenvalue.

is studied in [IW05a, IW05b, IW07, KW06, KW07, KY09]. Furthermore it is also known that the set of *odd* eigenvectors of non-commutative harmonic oscillator is deeply related to the set of some solutions of the Heun differential equation [IW05b, Och01, Och04]:

$$\frac{\partial^2}{\partial w^2}f+\left(\frac{1-n}{w}+\frac{-n}{w-1}+\frac{n+3/2}{w-a}\right)\frac{\partial}{\partial w}f+\frac{-(3/2)nw-q}{w(w-1)(w-a)}f=0,$$

where $n\in\mathbb{N}\cup\{0\}$, $a\in\mathbb{C}$ with $|a|<1$ and $q\in\mathbb{C}$.

In this paper we concentrate on the study of the lowest eigenvalue λ_1 of Q. We set

$$E=\lambda_1 \tag{1.2}$$

and $p=-id/dx$. In particular we are interested in determining the dimension of $\mathrm{Ker}(Q-E)$. The eigenvector associated with the lowest eigenvalue is called the ground state. In the case of $\alpha=\beta$, as is mentioned above, Q can be diagonalized as $Q\cong\begin{pmatrix}h&0\\0&h\end{pmatrix}$ with

$$h=\frac{1}{2}p^2+\frac{\alpha^2-1}{2}x^2, \tag{1.3}$$

where $\cong$ denotes the unitary equivalence. Then all the eigenvalues of $Q(\alpha,\alpha)$ are two-fold degenerate. In particular, its lowest eigenvalue

$$E_0=\frac{1}{2}\sqrt{\alpha^2-1} \tag{1.4}$$

is two-fold degenerate. In the general case, $\alpha\neq\beta$, the so-called Ichinose-Wakayama bound is established in [IW07]:

$$\left(j-\frac{1}{2}\right)\min\{\alpha,\beta\}\sqrt{\frac{\alpha\beta-1}{\alpha\beta}}\le\lambda_{2j-1}\le\lambda_{2j}\le\left(j-\frac{1}{2}\right)\max\{\alpha,\beta\}\sqrt{\frac{\alpha\beta-1}{\alpha\beta}} \tag{1.5}$$

for $j\in\mathbb{N}$. By this inequality we see that the multiplicity of E is at most two if $\beta<3\alpha$ or $\alpha<3\beta$.

Furthermore beyond above results one may expect that E is simple for $\alpha\neq\beta$. In [NNW02] it can be numerically shown that E is simple for $\alpha\neq\beta$ and, in [Par04] the simplicity is proven but only for sufficiently large $\alpha\beta$. It is then mentioned in [Par08b, 8.3 Notes] that the determining the multiplicity of the lowest eigenvalue should be explored.

In this paper we show that

- **(a):** E is at most twofold degenerate for $(\alpha,\beta)\in(2,\infty)\times(2,\infty)$,
- **(b):** E is simple for some region of α and β.

In order to prove (a), we apply the method in [Hir05], where the so-called pull-through formula [GJ68] is a key. The second result (b) consists of two estimates. The first is for large $|\beta-\alpha|$ and the second for small $|\beta-\alpha|$ but $\alpha\neq\beta$. The first case is proven in a similar manner to (a) and the second by the regular perturbation theory of discrete spectrum.

Let $\mathscr{G} = \mathrm{Ker}(Q - E)$ be the set of ground states. Let $L_+ \subset L^2(\mathbb{R})$ (resp. L_-) be the set of even functions (resp.odd functions). We define $\mathcal{H}_\pm = \mathbb{C}^2 \otimes L_\pm$. Since Q conserves the parity, Q is reduced by $\mathcal{H}_\pm$. Set $Q\lceil_{\mathcal{H}_\pm} = Q_\pm$ and then $Q = Q_+ \oplus Q_-$.

Theorem 1.1. *Suppose that* $(\alpha, \beta) \in (2, \infty) \times (2, \infty)$. *Then* $\dim\mathscr{G} \le 2$ *and* $\mathscr{G} \subset \mathcal{H}_+$. *I.e., the multiplicity of* E *is at most two and ground states are even functions.*

We can furthermore show that E is simple.

Theorem 1.2. *Suppose* $\beta > \alpha > 2$ *and*

$$\frac{1}{2} > \left(\frac{1}{2}\beta - E\right)^{-2} \frac{1}{\alpha^2 - 1} + \frac{1}{\alpha^2 - 1}. \tag{1.6}$$

Then E *is simple.*

Condition (1.6) includes the implicit value E. Let

$$E_{\mathrm{upper}} = \frac{\sqrt{\alpha\beta}\sqrt{\alpha\beta - 1}}{\alpha + \beta + |\alpha - \beta| \frac{(\alpha\beta-1)^{1/4}}{\sqrt{\alpha\beta}} \mathrm{Re}\rho},$$

where $\rho^2 = \sqrt{\alpha\beta - 1} - i$ with $\mathrm{Re}\,\omega > 0$, i.e., $\mathrm{Re}\,\rho = \sqrt{\dfrac{\sqrt{\alpha\beta}(\sqrt{\alpha\beta - 1} + 1)}{2}}$. Bound $E < E_{\mathrm{upper}}$ holds. See [Par08b, Theorem 8.2.1]. Combining this with (1.6) we have the corollary:

Corollary 1.3. *Suppose* $\beta > \alpha > 2$ *and*

$$\frac{1}{2} > \left(\frac{1}{2}\beta - E_{\mathrm{upper}}\right)^{-2} \frac{1}{\alpha^2 - 1} + \frac{1}{\alpha^2 - 1}. \tag{1.7}$$

Then E *is simple.*

Theorem 1.2 does not valid for (α, β) in a neighborhood of the diagonal line on $\alpha - \beta$ plane. See Figure 1. We can also however show that E is simple for α and β in a



FIGURE 1. The region satisfying (1.7)
(The case of $\alpha > \beta$ is also drawn.)

neighborhood of the diagonal line. We define $g_1, ..., g_4$ by

$$g_1 = (\alpha - 1)^{-1}\left(3 + \frac{\sqrt{3}}{\sqrt{\alpha^2-1}}\right), \tag{1.8}$$

$$g_2 = \frac{\sqrt{\alpha^2-1}}{2\left|\sqrt{\alpha^2-1} - \lambda_2\right|} g_1^2 \tag{1.9}$$

$$g_3 = \frac{\alpha}{2\sqrt{\alpha^2-1}} \tag{1.10}$$

$$g_4 = \frac{(\sqrt{\alpha^2-1})^{3/2}}{4\alpha^{3/2}}. \tag{1.11}$$

Theorem 1.4. *Let* $\varepsilon = \beta - \alpha$. *Assume that* $\beta > \alpha > 1$, $\sqrt{\beta^2-1} \le 3\sqrt{\alpha^2-1}$ *and* $\varepsilon^2 g_2 < 1/2$. *Let*

$$\kappa(\varepsilon) = E_0 g_1^2 + \varepsilon g_2(E_0 g_1 + g_3 + g_4) + \varepsilon^2 2E_0 g_1^2 g_2 + \varepsilon^3 2E_0 g_1 g_2^2, \tag{1.12}$$

$$\ell(\varepsilon) = (1 - \varepsilon^2 g_2)\sqrt{1 - 2\varepsilon^2 g_2^2}. \tag{1.13}$$

Then

$$|\lambda_1 - \lambda_2| \ge \frac{2\varepsilon}{\ell(\varepsilon)}(g_4 - \varepsilon\kappa(\varepsilon)) \tag{1.14}$$



FIGURE 2. A wedge-shaped region included in the region satisfying $\varepsilon\kappa(\varepsilon) < g_4$

In particular when $\varepsilon\kappa(\varepsilon) < g_4$, E is simple.

Note that we know the bound $\lambda_2 \leq \frac{\beta}{2}\frac{\sqrt{\alpha\beta-1}}{\sqrt{\alpha\beta}}$ by the Ichinose-Wakayama bound. Then, the region of α, β satisfying $\varepsilon\kappa(\varepsilon) < g_4$, includes a wedge-shaped region illustrated in Figure 1, where we also drew the case of $\alpha > \beta$.

2. PROOFS OF THEOREMS

2.1. **Proof of Theorem 1.1.** In the following we omit the symbol $\otimes$ for the notational simplicity, and we can suppose that $\alpha < \beta$ without loss of generality. Let

$$N = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}(p^2 + x^2 - 1).$$

The spectrum of N is $\sigma(N) = \{0\} \cup \mathbb{N}$ and the multiplicity of each eigenvalues. Let $a = \dfrac{1}{\sqrt{2}}(x + ip)$ and $a^* = \dfrac{1}{\sqrt{2}}(x - ip)$. They satisfy canonical commutation relations $[a, a^*] = 1$ and $[a, a] = [a^*, a^*] = 0$, and we have $a^*a = N$

Proof of Theorem 1.1: In terms of a and a^*, the operator Q can be realized as

$$Q = a^*Aa + \frac{1}{2}A + \frac{1}{2}\left(aJa - a^*Ja\right). \tag{2.1}$$

Let $\varphi_{\rm g} \in \mathscr{G}$. We have $(Q - E)a\varphi_{\rm g} = [Q, a]\varphi_{\rm g}$. Since $[Q, a] = -Aa + Ja^*$ by canonical commutation relation, we have $(Q - E)a\varphi_{\rm g} = (-Aa + Ja^*)\varphi_{\rm g}$ and then $(Q - E + A)a\varphi_{\rm g} =$

$Ja^*\varphi_{\rm g}$. Notice that $Q-E+A\geq\alpha>0$. We have

$$a\varphi_{\rm g}=(Q-E+A)^{-1}Ja^*\varphi_{\rm g}. \tag{2.2}$$

Taking the norm on both sides above, we have

$$\|a\varphi_{\rm g}\|^2\leq\frac{1}{\alpha^2}\|a^*\varphi_{\rm g}\|^2. \tag{2.3}$$

Since $\|a\varphi_{\rm g}\|^2=\langle\varphi_{\rm g},N\varphi_{\rm g}\rangle$ and $\|a^*\Omega\|=\langle\varphi_{\rm g},N\varphi_{\rm g}\rangle+\|\varphi_{\rm g}\|^2$, we see that

$$\langle\varphi_{\rm g},N\varphi_{\rm g}\rangle\leq\frac{1}{\alpha^2-1}\|\varphi_{\rm g}\|^2. \tag{2.4}$$

Let P_Ω be the projection onto $\ker N=\ker a$. Note that $N+\mathrm{P}_\Omega\geq 1$. Let $\mathscr{G}=\mathscr{G}_+\oplus\mathscr{G}_-$, where $\mathscr{G}_\pm=\mathscr{G}\cap\mathcal{H}_\pm$. Let $\mathrm{P}^\pm$ be the projection onto $\mathscr{G}_\pm$. Then, by (2.4), we have

$$\mathrm{P}^+\mathrm{P}_\Omega\mathrm{P}^+\geq\mathrm{P}^+(1-N)\mathrm{P}^+\geq\left(1-\frac{1}{\alpha^2-1}\right)\mathrm{P}^+. \tag{2.5}$$

Taking the trace of both sides, we have

$$2\geq\mathrm{Tr}(\mathrm{P}^+\mathrm{P}_\Omega\mathrm{P}^+)\geq\frac{\alpha^2-2}{\alpha^2-1}\mathrm{Tr}\mathrm{P}^+ \tag{2.6}$$

Thus we have the bound

$$\dim\ker\mathrm{P}^+\leq 2\frac{\alpha^2-1}{\alpha^2-2}. \tag{2.7}$$

Then the right hand side above is less than three for $\alpha>2$. Then $\dim\mathscr{G}^+\leq 2$. Similarly but replacing P^+ with P^-, we can also see that

$$\mathrm{P}^-\mathrm{P}_\Omega\mathrm{P}^-\geq\mathrm{P}^-(1-N)\mathrm{P}^-\geq\left(1-\frac{1}{\alpha^2-1}\right)\mathrm{P}^-. \tag{2.8}$$

Note that $\mathrm{P}^-\mathrm{P}_\Omega\mathrm{P}^-=0$, since P_Ω is the projection to the set of even functions. Then we have

$$0=\mathrm{Tr}(\mathrm{P}^-\mathrm{P}_\Omega\mathrm{P}^-)\geq\frac{\alpha^2-2}{\alpha^2-1}\mathrm{Tr}\mathrm{P}^-. \tag{2.9}$$

In particular for $\alpha>\sqrt{2}$, the dimension of $\mathscr{G}_-$ equal to zero. Then the theorem follows. □

2.2. **Proof of Theorem 1.2.** In this subsection we show that the lowest eigenvalue is simple. The strategy is parallel with that of previous subsection but P^+ is replaced by a projection R with the dimension of $\mathrm{Ran}\mathrm{R}=1$. Let $\sigma_1,\sigma_2,\sigma_3$ be the 2×2 Pauli matrices given by

$$\sigma_1=\begin{pmatrix}0&1\\1&0\end{pmatrix},\quad\sigma_2=\begin{pmatrix}0&-i\\i&0\end{pmatrix},\quad\sigma_3=\begin{pmatrix}1&0\\0&-1\end{pmatrix}. \tag{2.10}$$

Proof of Theorem 1.2: The Hamiltonian Q can be written in the form:

$$Q = \frac{1}{2}A(p^2 + x^2) + \frac{1}{2}\sigma_2(px + xp). \tag{2.11}$$

We set $M = \frac{1}{2}(1+\sigma_3) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $M^\perp = \frac{1}{2}(1-\sigma_3) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ and $\mathrm{R} = M^\perp \mathrm{P}_\Omega$. Then we have

$$(Q - E)\mathrm{R}\varphi_{\mathrm{g}} = M^\perp[Q, \mathrm{P}_\Omega]\varphi_{\mathrm{g}} + [Q, M^\perp]\mathrm{P}_\Omega\varphi_{\mathrm{g}}. \tag{2.12}$$

The commutator $[Q, M^\perp]$ can be computed as

$$[Q, M^\perp] = -\frac{1}{2}(aa - a^*a^*)\sigma_1. \tag{2.13}$$

Thus we have

$$[Q, \mathrm{R}] = M^\perp \frac{i}{2}\sigma_2[aa - a^*a^*, \mathrm{P}_\Omega] + \frac{1}{2}a^*a^*\sigma_1\mathrm{P}_\Omega, \tag{2.14}$$

where we used the fact that $a\mathrm{P}_\Omega = 0$. Hence we have

$$\begin{aligned} \langle \mathrm{R}\varphi_{\mathrm{g}}, (Q-E)\mathrm{R}\varphi_{\mathrm{g}}\rangle &= \left\langle \varphi_{\mathrm{g}}, \frac{i}{2}\mathrm{R}M^\perp\sigma_2[aa - a^*a^*, \mathrm{P}_\Omega]\varphi_{\mathrm{g}} \right\rangle + \frac{1}{2}\left\langle \varphi_{\mathrm{g}}, \mathrm{R}a^*a^*\sigma_1\mathrm{P}_\Omega\varphi_{\mathrm{g}}\right\rangle \\ &= \frac{i}{2}\left\langle \varphi_{\mathrm{g}}, \mathrm{R}\sigma_2 a^2 \varphi_{\mathrm{g}}\right\rangle. \end{aligned} \tag{2.15}$$

On the other hand $\mathrm{R}(Q-E)\mathrm{R} = \left(\frac{1}{2}\beta - E\right)\mathrm{R}^2$. Then (2.4) and $\|a^*\mathrm{R}\varphi_{\mathrm{g}}\| = \|\mathrm{R}\varphi_{\mathrm{g}}\|$ yield that

$$\left(\frac{1}{2}\beta - E\right)\|\mathrm{R}\varphi_{\mathrm{g}}\|^2 \le \frac{1}{2}\|a^*\mathrm{R}\varphi_{\mathrm{g}}\|\|\sigma_2 a\varphi_{\mathrm{g}}\| \le \frac{1}{\sqrt{\alpha^2-1}}\|\mathrm{R}\varphi_{\mathrm{g}}\|\|\varphi_{\mathrm{g}}\|.$$

Therefore

$$\|\mathrm{R}\varphi_{\mathrm{g}}\|^2 \le \left(\frac{1}{2}\beta - E\right)^{-2}\frac{1}{\alpha^2-1}\|\varphi_{\mathrm{g}}\|^2.$$

Since $M + M^\perp = 1$, it holds that $\mathrm{P}_\Omega M + \mathrm{R} + N \ge 1$. Then, by using (2.4) we have

$$\mathrm{P}(\mathrm{P}_\Omega M)\mathrm{P} \ge \mathrm{P}(1 - \mathrm{P}_\Omega M^\perp - N)\mathrm{P} = \left(1 - \left(\frac{1}{2}\beta - E\right)^{-2}\frac{1}{\alpha^2-1} - \frac{1}{\alpha^2-1}\right)\mathrm{P}$$

where $P = P^+ + P^-$ is the orthogonal projection onto $\mathscr{G}$. Taking the trace of both sides above, we have

$$1 \ge \left(1 - \left(\frac{1}{2}\beta - E\right)^{-2}\frac{1}{\alpha^2-1} - \frac{1}{\alpha^2-1}\right)\mathrm{Tr}\mathrm{P},$$

and the theorem follows. □

2.3. **Proof of Theorem 1.4.** Recall that $\varepsilon = \beta - \alpha$. In this section, we fix an arbitrary $\alpha > 1$ and set

$$Q(\alpha, \beta) = Q = Q(\varepsilon) = Q_0 + \varepsilon V, \tag{2.16}$$

where $Q_0 = Q(\alpha, \alpha)$ and $V = \frac{1}{2}\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}(p^2 + x^2)$.

Lemma 2.1. *For all $\Phi \in D(Q_0)$, it follows that*

$$\|V\Phi\| \le (\alpha - 1)^{-1}\|Q_0\Phi\| + \frac{\sqrt{3}}{2}(\alpha - 1)^{-1}\|\Phi\|. \tag{2.17}$$

Proof. One can show that $\|(px + xp)u\|^2 \le \|(p^2 + x^2)u\|^2 + 3\|u\|^2$. Since $\sigma_2^2 = 1$, we have

$$\begin{aligned} \|Q_0\Phi\| \ge & \alpha\|\frac{1}{2}(p^2 + x^2)\Phi\| - \|\frac{1}{2}(px + xp)\Phi\| \\ \ge & \alpha\|\frac{1}{2}(p^2 + x^2)\Phi\| - \frac{1}{2}\left(\|(p^2 + x^2)\Phi\|^2 + 3\|\Phi\|^2\right)^{1/2}, \end{aligned}$$

and hence

$$\|Q_0\Phi\| \ge \frac{\alpha - 1}{2}\|(p^2 + x^2)\Phi\| - \frac{\sqrt{3}}{2}\|\Phi\|. \tag{2.18}$$

Noticing $\|V\Phi\| \le \frac{1}{2}\|(p^2 + x^2)\Phi\|$ we have the bound (2.17). □

By (1.5) or the sandwich estimate $Q(\alpha, \alpha) \le Q(\alpha, \beta) \le Q(\beta, \beta)$ we see bounds: $\frac{\sqrt{\alpha^2-1}}{2} \le \lambda_1 \le \lambda_2 \le \frac{\sqrt{\beta^2-1}}{2}$ and $\frac{3}{2}\sqrt{\alpha^2 - 1} \le \lambda_3$. When $\sqrt{\beta^2 - 1} \le 3\sqrt{\alpha^2 - 1}$, Q has exactly two eigenvalues in the interval $\left[\frac{1}{2}\sqrt{\alpha^2 - 1}, \frac{1}{2}\sqrt{\beta^2 - 1}\right]$. Let C be the closed disk centered at $\frac{\sqrt{\alpha^2-1}}{2}$ with the radius $\frac{\sqrt{\alpha^2-1}}{2}$ in the complex plane:

$$C = \left\{ \frac{\sqrt{\alpha^2 - 1}}{2} + re^{i\theta} \in \mathbb{C} \,\middle|\, 0 \le r \le \frac{\sqrt{\alpha^2 - 1}}{2}, 0 \le \theta \le 2\pi \right\}. \tag{2.19}$$

Thus Q has exactly two eigenvalues, λ_1 and λ_2, inside of C.

Lemma 2.2. *It follows that $\|V(Q_0 - z)^{-1}\| \le g_1$ for all $z \in \partial C$.*

Proof. By Lemma 2.1, we have

$$\|V(Q_0 - z)^{-1}\| \le (\alpha - 1)^{-1}\|Q_0(Q_0 - z)^{-1}\| + \frac{\sqrt{3}}{2}(\alpha - 1)^{-1}\|(Q_0 - z)^{-1}\|. \tag{2.20}$$

Since the eigenvalues of Q_0 are $\{(\frac{1}{2} + n)\sqrt{\alpha^2 - 1}\}_{n=0}^{\infty}$ we have $\sup_{z \in \partial C}\|Q_0(Q_0 - z)^{-1}\| = 3$ and $\sup_{z \in \partial C}\|(Q_0 - z)^{-1}\| = \frac{2}{\sqrt{\alpha^2-1}}$. Then the lemma follows. □

The two-dimensional subspace spanned by eigenvectors associated with eigenvalues λ_1 and λ_2 is denoted by $\mathscr{F}$. The orthogonal projection onto $\mathscr{F}$ is then given by

$$P = P(\varepsilon) = -\frac{1}{2\pi i} \oint_{\partial C} (Q - z)^{-1} dz. \tag{2.21}$$

We expand $P(\varepsilon)$ with respect to ε up to the second order:

$$P = P_0 + \varepsilon P_1 + \varepsilon^2 R, \tag{2.22}$$

where P_0 is the orthogonal projection onto the ground states of Q_0 and

$$P_1 = -\frac{1}{2\pi i} \oint_{\partial C} (Q_0 - z)^{-1} V (Q_0 - z)^{-1} dz, \tag{2.23}$$

$$R = R(\varepsilon) = -\frac{1}{2\pi i} \oint_{\partial C} (Q_0 - z)^{-1} V (Q_0 - z)^{-1} V (Q - z)^{-1} dz. \tag{2.24}$$

Lemma 2.3. *We have* $\|R\| \le g_2$ *and* $\|VP_1\| \le E_0 g_1^2$*.*

Proof. By Lemma 2.2, we have

$$\|R\| \le \frac{|C|}{2\pi} \sup_{z\in\partial C} \|V(Q_0 - z)^{-1}\|^2 \|(Q - z)^{-1}\| \le \frac{\sqrt{\alpha^2 - 1}}{2} g_1^2 \|(Q - \sqrt{\alpha^2 - 1})^{-1}\|. \tag{2.25}$$

Since $\lambda_3 \ge \frac{3}{2}\sqrt{\alpha^2 - 1}$, we have $\|(Q - \sqrt{\alpha^2 - 1})^{-1}\| = |\lambda_2 - \sqrt{\alpha^2 - 1}|^{-1}$. Hence $\|R\| \le g_2$ holds. Similarly one can prove the second bound. $\square$

Let $v_0 \in L^2(\mathbb{R})$ be the normalized ground state of $h = \frac{1}{2}p^2 + \frac{\alpha^2-1}{2}x^2$. Namely

$$v_0(x) = \left(\frac{\sqrt{\alpha^2 - 1}}{\pi}\right)^{1/4} e^{-\sqrt{\alpha^2-1}x^2/2}. \tag{2.26}$$

Let S_a be the dilation defined by $S_a f(x) = \frac{1}{\sqrt{a}} f(ax)$ for $a > 0$. We define the unitary operator U on $\mathcal{H}$ by

$$U = \frac{1}{\sqrt{2}} S_{\sqrt{\alpha}} \begin{pmatrix} e^{ix^2/(2\alpha)} & 0 \\ 0 & e^{-ix^2/(2\alpha)} \end{pmatrix} \begin{pmatrix} 1 & -i \\ 1 & i \end{pmatrix}. \tag{2.27}$$

Then $UQ_0U^* = \begin{pmatrix} h & 0 \\ 0 & h \end{pmatrix}$ and vectors $u_1 = U^* \begin{pmatrix} v_0 \\ 0 \end{pmatrix}$ and $u_2 = U^* \begin{pmatrix} 0 \\ v_0 \end{pmatrix}$ are two fold ground states of $Q(\alpha, \alpha)$. Since P is a projection onto $\mathscr{F}$, each of the ground state and the first excited state can be expresses as a linear combination of Pu_1 and Pu_2 as long as both Pu_1 and Pu_2 are linearly independent, which is proven in the lemma below:

Lemma 2.4. *Assume that* $\varepsilon^2 g_2 < 1$*. Then* Pu_1 *and* Pu_2 *are non-zero vectors. Moreover, if* $\varepsilon^2 g_2 < 1/2$*, then* Pu_1 *and* Pu_2 *are linearly independent.*

Proof. By (2.22) we have

$$\|Pu_1\|^2 = 1 + \varepsilon \langle u_1, P_1 u_1 \rangle + \varepsilon^2 \langle u_1, R u_1 \rangle . \tag{2.28}$$

The second term on the right hand side above is zero, since

$$\langle u_1, P_1 u_1 \rangle = \frac{-1}{2\pi i} \oint_{\partial C} \frac{1}{(E_0 - z)^2} dz \langle u_1, V u_1 \rangle = 0. \tag{2.29}$$

Hence $\|Pu_1\|^2 \geq 1 - \varepsilon^2 g_2^2 > 0$ holds by Lemma 2.3 and the assumption. Thus $Pu_i (i = 1, 2)$ are non-zero vectors. Next we assume that $\varepsilon^2 g_2 < 1/2$. Then we have

$$\begin{aligned} \|Pu_1\|^2 \|Pu_2\|^2 - | \langle Pu_1, Pu_2 \rangle |^2 &\geq (1 - \varepsilon^2 g_2)(1 - \varepsilon^2 g_2) - \varepsilon^4 g_2^2 \\ &= 1 - 2\varepsilon^2 g_2 > 0 \end{aligned} \tag{2.30}$$

which implies that Pu_1 and Pu_2 are linearly independent. □

Lemma 2.5. *Let g_3 and g_4 be given in (1.10) and (1.11), respectively. Then it follows that where we used the bound*

$$\|Vu_i\| \leq g_1 E_0, \quad i = 1, 2, \tag{2.31}$$

$$| \langle u_1, V u_1 \rangle | = g_3, \tag{2.32}$$

$$| \langle u_1, V u_2 \rangle | \leq g_4. \tag{2.33}$$

Proof. (2.31) follows from

$$\|Vu_i\| \leq \|V(Q_0 - \sqrt{\alpha^2 - 1})^{-1}\| \|(Q_0 - \sqrt{\alpha^2 - 1}) u_i\| \leq g_1 E_0. \tag{2.34}$$

The proofs of (2.32) and (2.33) are given in Appendix. □

Proof of Theorem 1.4: Suppose that $\varepsilon^2 g_2 < 1/2$. We define

$$\Phi_1 = \frac{Pu_1}{\|Pu_1\|}, \qquad \Phi_2 = \frac{Pu_2 - \langle \Phi_1, u_2 \rangle \Phi_1}{\|Pu_2 - \langle \Phi_1, u_2 \rangle \Phi_1\|} \tag{2.35}$$

Then, by Lemma 2.4, Φ_1 and Φ_2 are orthogonal vectors in $\mathscr{F}$. Let $\mathscr{V} = L.H.\{\Phi_1, \Phi_2\}$ be the two dimensional vector space and $Q : \mathscr{V} \to \mathscr{V}$ can be regarded as a linear operator and its matrix representation is given by

$$m = \begin{pmatrix} \langle \Phi_1, Q\Phi_1 \rangle & \langle \Phi_1, Q\Phi_2 \rangle \\ \langle \Phi_2, Q\Phi_1 \rangle & \langle \Phi_2, Q\Phi_2 \rangle \end{pmatrix} . \tag{2.36}$$

Thus the eigenvalues λ_1 and λ_2 of Q are also the eigenvalue of m. Therefore, the difference of λ_1 and λ_2 can be computed by

$$(\lambda_2 - \lambda_1)^2 = (\langle \Phi_1, Q\Phi_1 \rangle - \langle \Phi_2, Q\Phi_2 \rangle)^2 + 4 | \langle \Phi_1, Q\Phi_2 \rangle |^2, \tag{2.37}$$

which implies that

$$|\lambda_2 - \lambda_1| \geq 2 | \langle \Phi_1, Q\Phi_2 \rangle |. \tag{2.38}$$

We estimate $|\langle\Phi_1, Q\Phi_2\rangle|$ from below. Inserting the definition of Φ_j into $\langle\Phi_i, Q\Phi_j\rangle$ we have

$$\langle\Phi_1, Q\Phi_2\rangle = \frac{\langle Pu_1, Q(\|Pu_1\|^2 Pu_2 - (Pu_1, Pu_2)Pu_1)\rangle}{\|Pu_1\|^3 \|Pu_2 - (Pu_1, Pu_2)Pu_1/\|Pu_1\|^2\|}. \tag{2.39}$$

Notice that

$$\langle Pu_i, QPu_j\rangle = (Pu_i, Qu_j) = E_0\delta_{ij} + \varepsilon\langle V\rangle_{ij} + \varepsilon^2(E_0\langle R\rangle_{ij} + \langle P_1V\rangle_{ij}) + \varepsilon^3\langle RV\rangle_{ij},$$

where $\langle K\rangle_{ij} = (u_i, Ku_j)$. We see that the denominator of (2.39) is expanded as

$$\begin{aligned}&\|Pu_1\|\|\|Pu_1\|^2 Pu_2 - (Pu_1, Pu_2)Pu_1\|\\ &= (1+\varepsilon^2\langle R\rangle_{11})\sqrt{(1+\varepsilon^2\langle R\rangle_{11})(1+\varepsilon^2\langle R\rangle_{22}) - \varepsilon^4|\langle R\rangle_{12}|^2}.\end{aligned}$$

By the bound $\|R\| \le g_2$ we have the lower bound

$$\|Pu_1\|\|\|Pu_1\|^2 Pu_2 - (Pu_1, Pu_2)Pu_1\| \ge (1-\varepsilon^2 g_2)\sqrt{1-2\varepsilon^2 g_2^2}. \tag{2.40}$$

The numerator of (2.39) can be also expanded as

$$\begin{aligned}&(Pu_1, Q(\|Pu_1\|^2 Pu_2 - (Pu_1, Pu_2)Pu_1))\\ &= \varepsilon\langle V\rangle_{12} + \varepsilon^2\langle P_1V\rangle_{12} + \varepsilon^3\Big(\langle RV\rangle_{12} + \langle V\rangle_{12}\langle R\rangle_{11} - \langle R\rangle_{12}\langle V\rangle_{11}\Big)\\ &\quad + \varepsilon^4\Big(\langle P_1V\rangle_{12}\langle R\rangle_{11} - \langle R\rangle_{12}\langle P_1V\rangle_{11}\Big) + \varepsilon^5\Big(\langle R\rangle_{11}\langle RV\rangle_{12} - \langle R\rangle_{12}\langle RV\rangle_{12}\Big).\end{aligned}$$

By using Lemmas 2.3 and 2.5, each term can be evaluated as

$$\begin{aligned}&\varepsilon^2|\langle P_1V\rangle_{12}| \le \varepsilon^2 E_0 g_1^2\\ &\varepsilon^3|\langle RV\rangle_{12} + \langle V\rangle_{12}\langle R\rangle_{11} - \langle R\rangle_{12}\langle V\rangle_{11}| \le \varepsilon^3\left(E_0 g_1 g_2 + g_2 g_4 + g_2 g_3\right)\\ &\varepsilon^4|\langle P_1V\rangle_{12}\langle R\rangle_{11} - \langle R\rangle_{12}\langle P_1V\rangle_{11}| \le \varepsilon^4 2E_0 g_1^2 g_2\\ &\varepsilon^5|\langle R\rangle_{11}\langle RV\rangle_{12} - \langle R\rangle_{12}\langle RV\rangle_{12}| \le \varepsilon^5 2E_0 g_1 g_2^2.\end{aligned}$$

By combining all the estimates stated above, we have

$$|\lambda_1 - \lambda_2| \ge \frac{2\varepsilon}{\ell(\varepsilon)}\left(g_4 - \varepsilon\kappa(\varepsilon)\right), \tag{2.41}$$

where $\ell(\varepsilon) = (1-\varepsilon^2 g_2)\sqrt{1-2\varepsilon^2 g_2^2}$. Hence the theorem follows. □

Appendix A. Computation of $\langle V\rangle_{ij}$

We recall that $v_0=(\omega/\pi)^{1/4}e^{-\omega x^2/2}$ with $\omega=\sqrt{\alpha^2-1}$, $V=\frac{1}{2}\begin{pmatrix}0&0\\0&1\end{pmatrix}(p^2+x^2)$ and U is given by (2.27). We directly see that

$$
\begin{aligned}
&UVU^{-1}\\
&=\frac{1}{4}S_{\sqrt{\alpha}}\begin{pmatrix}e^{ix^2/2\alpha}&0\\0&e^{-ix^2/2\alpha}\end{pmatrix}\begin{pmatrix}1&-i\\1&i\end{pmatrix}\begin{pmatrix}0&0\\0&1\end{pmatrix}\begin{pmatrix}1&1\\i&-i\end{pmatrix}(p^2+x^2)\begin{pmatrix}e^{-ix^2/2\alpha}&0\\0&e^{ix^2/2\alpha}\end{pmatrix}S_{1/\sqrt{\alpha}}\\
&=\frac{1}{4}\begin{pmatrix}e^{ix^2/2\alpha}(\frac{p^2}{\alpha}+\alpha x^2)e^{-ix^2/2\alpha}&e^{-ix^2/2\alpha}(\frac{p^2}{\alpha}+\alpha x^2)e^{ix^2/2\alpha}\\-e^{-ix^2/2\alpha}(\frac{p^2}{\alpha}+\alpha x^2)e^{-ix^2/2\alpha}&e^{ix^2/2\alpha}(\frac{p^2}{\alpha}+\alpha x^2)e^{ix^2/2\alpha}\end{pmatrix}.
\end{aligned}
$$

Then we have

$$
\left\langle\begin{pmatrix}1\\0\end{pmatrix},UVU^{-1}\begin{pmatrix}1\\0\end{pmatrix}\right\rangle_{\mathbb{C}^2}=\frac{1}{4}e^{ix^2/2\alpha}\left(\frac{p^2}{\alpha}+\alpha x^2\right)e^{-ix^2/2\alpha} \tag{1.1}
$$

$$
\left\langle\begin{pmatrix}1\\0\end{pmatrix},UVU^{-1}\begin{pmatrix}0\\1\end{pmatrix}\right\rangle_{\mathbb{C}^2}=-\frac{1}{4}e^{ix^2/2\alpha}\left(\frac{p^2}{\alpha}+\alpha x^2\right)e^{ix^2/2\alpha}. \tag{1.2}
$$

Lemma A.1. *It follows that* $\langle u_1,Vu_1\rangle=\dfrac{\alpha}{4\omega}$, *and then* $|\langle u_1,Vu_1\rangle|=g_3$.

Proof. By (1.1) we have

$$
\langle u_1,Vu_1\rangle=\frac{1}{4}\left\langle v_0,e^{ix^2/2}\left(\frac{p^2}{\alpha}+\alpha x^2\right)e^{-ix^2/2}v_0\right\rangle=\frac{1}{4}\left\langle v_0,\left(\frac{(p-x)^2}{\alpha}+\alpha x^2\right)v_0\right\rangle.
$$

Since $\langle v_0,(px+xp)v_0\rangle=0$, we obtain that

$$
\langle u_1,Vu_1\rangle=\frac{1}{4}\left\langle v_0,\left(\frac{p^2+x^2}{\alpha}+\alpha x^2\right)v_0\right\rangle.
$$

By $\langle v_0,x^2v_0\rangle=1/(2\omega)$ and $\langle v_0,p^2v_0\rangle=\omega/2$ we have $\langle u_1,Vu_1\rangle=\alpha/(4\omega)$. □

Lemma A.2. *It follows that* $\langle u_1,Vu_2\rangle=-\dfrac{\omega^{3/2}}{4\alpha}(\omega-i)^{-1/2}$. *In particular* $|\langle u_1,Vu_2\rangle|\le\dfrac{\omega^{3/2}}{4\alpha^{3/2}}$.

Proof. By (1.2) we have

$$
\begin{aligned}
\langle u_1,Vu_2\rangle&=-\frac{1}{4}\left\langle v_0,e^{ix^2/2}\left(\frac{p^2}{\alpha}+\alpha x^2\right)e^{ix^2/2}v_0\right\rangle\\
&=-\frac{1}{4\alpha}\left\langle(-i-\omega)xe^{-ix^2/2}v_0,(i-\omega)xe^{ix^2/2}v_0\right\rangle-\frac{\alpha}{4}\left\langle v_0,x^2e^{ix^2}v_0\right\rangle\\
&=-\frac{1}{4\alpha}((\omega-i)^2+\alpha^2)\left\langle v_0,x^2e^{ix^2}v_0\right\rangle.
\end{aligned}
$$

Since $\left\langle v_0, x^2 e^{ix^2} v_0 \right\rangle = (\omega/\pi)^{1/2} \int_{\mathbb{R}} x^2 e^{-(\omega-i)x^2} dx = \dfrac{1}{2}\omega^{1/2}(\omega-i)^{-3/2}$, we have the lemma.

$\square$

Acknowledgment

We thank Takashi Ichinose and Masato Wakayama for sending [IW07] to our attention and giving helpful comments. I. S.'s work was partially supported by Y22740087 from JSPS, and was performed through the Program for Dissemination of Tenure-Track System funded by the Ministry of Education and Science, Japan. F.H. thanks for the financial support by Grant-in-Aid for Science Research (B) 20340032 from JSPS.

References

[GJ68] J. Glimm and A. Jaffe, The $\lambda(\phi^4)_2$ quantum field theory without cutoffs. I. *Phys. Rev.* **176** (1968), 1945–1951.

[HH12] M. Hirokawa and F. Hiroshima, Absence of energy level crossing for the ground state energy of the Rabi model, preprint 2012.

[Hir05] F. Hiroshima, Multiplicity of ground states in quantum field models:applications of asymptotic fields, *J. Funct. Anal.* **224** (2005), 431–470.

[IW05a] T. Ichinose and M. Wakayama, Zeta functions for the spectrum of the non-commutative harmonic oscillators, *Commun. Math. Phys.* **258** (2005), 697–739.

[IW05b] T. Ichinose and M. Wakayama, Special values of the spectral zeta function of the non-commutative harmonic oscillator and confluent Heun equations, *Kyushu. J. Math.* **59** (2005), 39–100.

[IW07] T. Ichinose and M. Wakayama, On the spectral zeta function for the non-commutative harmonic oscillator and, *Rep. Math. Phys.* **59** (2007), 421–432.

[KW06] K. Kimoto and M. Wakayama, Apéry-like number arising from special values of spectral zeta functions for non-commutative harmonic oscillators, *Kyushu. J. Math.* **60** (2006), 383–404.

[KW07] K. Kimoto and M. Wakayama, Elliptic curves arising from the spectral zeta function for non-commutative harmonic oscillators and $\Gamma_0(4)$-modular forms, *The Conference on L-Functions*, World Sci. Publ., Hackensack, NJ, 2007, 201–218.

[KY09] K. Kimoto and Y. Yamasaki, A variation of multiple L-values arising from the spectral zeta function of the non-commutative harmonic oscillator, *Proc. Amer. Math. Soc.* **137** (2009), 2503–2515.

[NNW02] K. Nagatou, M. T. Nakao and M. Wakayama, Verified numerical computations for eigenvalues of non-commutative harmonic oscillators, *Num. Funct. Anal. and Opt.* **23** (2002), 633–650.

[Och01] H. Ochiai Non-commutative harmonic oscillators and Fuchsian ordinary differential operators, *Commun. Math. Phys.* **217** (2001), 357–373.

[Och04] H. Ochiai Non-commutative harmonic oscillators and the connection problem for the Heun differential equation, *Lett. Math. Phys.* **70** (2004), 133–139.

[PW01] A. Parmeggiani and M. Wakayama, Oscillator representations and systems of ordinary differential equations, *Proceedings of the National Academy of Sciences U.S.A* **98** (2001), 26–30.

[PW02a] A. Parmeggiani and M. Wakayama, Non-commutative harmonic oscillator I, *Form Math.* **14** (2002), 539–604.

[PW02b] A. Parmeggiani and M. Wakayama, Non-commutative harmonic oscillator II, *Form Math.* **14** (2002), 669–690.

[PW03] A. Parmeggiani and M. Wakayama, Corrigenda and remarks to: "Non-commutative harmonic oscillators. I." [Forum Math. 14 (2002), 539–604] *Forum Math.* **15** (2003), 955–963.

[Par04] A. Parmeggiani, On the spectrum and the lowest eigenvalue of certain non-commutative harmonic oscillators, *Kyushu J. Math.* **58** (2004), 277–322.

[Par06] A. Parmeggiani, On the spectrum of certain noncommutative harmonic oscillators, *Ann. Univ. Ferrara Sez. VII Sci. Mat.* **52** (2006), 431–456.

[Par08a] A. Parmeggiani, On the spectrum of certain non-commutative harmonic oscillators and semiclassical analysis, *Comm. Math. Phys.* **279** (2008), 285–308.

[Par08b] A. Parmeggiani, *Introduction to the spectral theory of non-commutative harmonic oscillators*, COE Lecture Note **8**, Kyushu University, 2008.

[Par10] A. Parmeggiani, *Spectral theory of non-commutative harmonic oscillators: an introduction*, Lecture Notes in Mathematics, **1992**, Springer-Verlag, Berlin, 2010.

Faculty of Mathematics, Kyushu University, Fukuoka, 819-0395, Japan
E-mail address: `hiroshima@math.kyushu-u.ac.jp.`

Fiber-Nanotech Young Researcher Empowerment Center, Shinshu University, Matsumoto 390–8621, Japan
E-mail address: `isasaki@shinshu-u.ac.jp`